



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

XVII. *Observation of the Winter Solstice of 1812, with the Mural Circle at Greenwich.* By John Pond, Esq. *Astronomer Royal, F. R. S.*

Read February 25, 1813.

THE weather this year at the period of the solstice was peculiarly unfavourable for astronomical observation; however, in the course of the month, I obtained nine observations of the sun; one of these proved defective, the result of the other eight, accompanies this communication. In my observation of the summer solstice, it will be seen that I assumed the arc ZP equal $38^{\circ} 31' 21'' 15$; by subsequent observation I conceive that I have somewhat improved this quantity, which I now assume $38^{\circ} 31' 21'' 5$, as resulting from 120 observations of Polaris.

The observation of the summer solstice thus corrected will give the mean obliquity of the ecliptic for January 1, 1813, $23^{\circ} 27' 51'' 50$, and the winter solstice $23^{\circ} 27' 47'' 35$.

There can be no doubt but this small discordance might easily be reconciled by a slight modification of BRADLEY's refractions, and perhaps ultimately it may be necessary to have recourse to this theory for its explanation; but I am unwilling to do this hastily, being now occupied in making an extensive series of observations of circumpolar stars, with a view of determining, if possible, whether BRADLEY's mean refraction does, or does not, require alteration.—As I propose making the discordance of the solstices the subject of a sepa-

rate paper, I shall not add any thing farther on the subject, excepting a recommendation to astronomers, to deduce their refractions from circumpolar stars altogether, and then examine the solstices with those refractions, and by no means to make the coincidence of the solstices a required proof of their accuracy.

Note. In the annexed computation the reduction to the solstice is computed from the *observed* right ascension of the sun, by this means the errors of the tables being avoided, observations at a considerable distance (even some weeks from the solstice) may be employed, particularly in the winter season, when the uncertainty from refraction is greater than the error which may be probably introduced into this part of the calculation.

1812.	Barometer:		Therm.	Refraction:		Observations as given by the Instrument.	Position of the Zero point, or Equation to be applied to obtain the N. P. D.	Equation to reduce the ob- served N. P. D. to Zenith Distance or position of the Zenith point on the Circle.	Semi-diameter of the ☉, greater than in the Nautical Almanack.	Reduction to the Solstice.	Solstitial Zenith Distance.		N. P. D.
	In.	Out.									° ' "	° ' "	
Dec. 6 30,19	41	38	3	21,1	° ' "	☉ UL 112 12 34,3	+ 0 6,5	38 31 15,0	+ 16 17,3	' 31,3	74 56 29,0	113 27 50,5	
7 30,42	34	29	3	35,2		LL 112 51 44,4	6,5	15,0	— 16 17,3	' 37,4	24,7	46,2	
9 29,81	29	24	3	36,5		LL 113 4 14,8	6,5	15,0	— 16 17,5	' 36 10,0	28,8	50,3	
10 29,73	33	30	3	27,6		UL 112 37 19,5	6,5	15,0	+ 16 17,6	' 30 36,8	26,5	48,0	
11 29,84	35	34	3	30,0		LL 113 14 49,1	6,5	15,0	— 16 17,7	' 25 30,3	22,7	44,2	
13 29,61	32	30	3	30,6		UL 112 51 13,0	6,5	15,0	+ 16 18,0	' 16 41,8	28,4	49,9	
15 29,44	32	31	3	39,3		LL 113 30 39,6	6,5	15,0	— 16 18,1	' 9 43,4	29,2	50,7	
30 30,10	45	47	3	26,8		UL 112 50 38,8	6,5	15,0	+ 16 18,8	' 17 18,0	27,4	48,9	
Mean of 8 Observations													
Nutation + 7",62 Parallax — 8",50 ☉ Lat. — 0",36 =													
											74 56 27,09	113 27 48,59	
											— 1,24	— 1,24	
											74 56 25,85	113 27 47,35	